

KOBE-FHD-97-01

FUT-TH-97-01

May 1997

Hard Processes with Spin-Dependent Parton Distributions

T. Yamanishi[†], T. Morii¹, D. Roy¹, K. Sudoh² and S. Tanaka³

Research Center for Nuclear Physics, Osaka University, Ibaraki, Osaka, 567, Japan

¹*Faculty of Human Development, Kobe University, Nada, Kobe 657, Japan*

²*Department of Education, Kobe University, Nada, Kobe 657, Japan*

³*Department of Physics, Hyogo University of Education, Yashiro-cyo, Hyogo 673-14, Japan*

ABSTRACT

To test a possible color-octet contribution of heavy quarkonium productions, we propose ψ' -productions for small- p_T regions, where the color-singlet ψ' state cannot be produced by the gluon-gluon fusion owing to charge conjugation. We also point out that these processes are very useful to deduce informations of spin-dependent gluon distributions Δg .

PACS numbers: 13.85.Ni, 13.88.+e, 14.20.Dh, 14.40.Lb, 14.70.Dj

Recently, it has been reported that the cross sections of heavy quarkonium productions measured in unpolarized $p\bar{p}$ collisions by the CDF collaboration are inconsistent with the calculation with the lowest order processes alone[1]. Several groups have proposed new mechanisms: the production proceeds with not only color-singlet but also color-octet quarkonium states and this new mechanism dominates over the lowest order processes for large- p_T quarkonium productions[2]. But it does not seem to work for $\gamma + p \rightarrow J/\psi + X$ [3]. The discussion on the color-octet contribution is still controversial. Thus, it is important to test this mechanism in other experiments.

On the other hand, measurements of the spin-dependent structure functions of a nucleon $g_1(x, Q^2)$ for polarized deep inelastic scatterings (PDIS) imply a surprising result that the amount of the proton spin carried by quarks, ($\sim 30\%$), is much smaller than expected

[†]E-mail: yamanisi@natura.h.kobe-u.ac.jp , Phone: +81-6-879-8940 , Fax: +81-6-879-8899 .

from the quark–parton model[4]. One of keys to solve this spin puzzle is to directly measure the spin–dependent gluon distributions Δg and study the behavior of them. Relativistic Heavy Ion Collider (RHIC) at Brookhaven National Laboratory (BNL), which will start early in the next decade, could offer such a possibility and will give us fruitful informations on the nucleon spin structure.

In this talk, to test a possible color–octet contribution to charmonium productions[2] and furthermore to deduce informations of the spin–dependent gluon distributions, we suggest ψ' –productions for small– p_T regions at RHIC energies, where the color–singlet ψ' state does not contribute to the process due to charge conjugation. This process proceeds via the gluon–gluon fusion in the lowest order. Although there are several charmonium states, the detection for ψ' is easier than that for other charmonium states.

Let us introduce a two–spin asymmetry A_{LL} for this process,

$$A_{LL} = \frac{[d\sigma_{++} - d\sigma_{+-} + d\sigma_{--} - d\sigma_{-+}]}{[d\sigma_{++} + d\sigma_{+-} + d\sigma_{--} + d\sigma_{-+}]} = \frac{d\Delta\sigma}{d\sigma}, \quad (1)$$

where $d\sigma_{+-}$, for instance, denotes that the helicity of one beam particle is positive and the other is negative.

The spin–dependent and spin–independent differential cross sections are

$$\begin{aligned} \frac{d\Delta\sigma}{dx_L} = & \frac{\tau}{\sqrt{x_L^2 + 4\tau}} \left[\frac{\pi^3 \alpha_s^2}{144m_c^5} \{ \langle \mathcal{O}_8^{\psi'}(^1S_0) \rangle - \frac{1}{m_c^2} \langle \mathcal{O}_8^{\psi'}(^3P_0) \rangle \} \Delta g(x_a, Q^2) \Delta g(x_b, Q^2) \right. \\ & \left. - \frac{\pi^3 \alpha_s^2}{54m_c^5} \langle \mathcal{O}_8^{\psi'}(^3S_1) \rangle \{ \Delta q(x_a, Q^2) \Delta \bar{q}(x_b, Q^2) + \Delta q \leftrightarrow \Delta \bar{q} \} \right], \quad (2) \end{aligned}$$

$$\begin{aligned} \frac{d\sigma}{dx_L} = & \frac{\tau}{\sqrt{x_L^2 + 4\tau}} \left[\frac{\pi^3 \alpha_s^2}{144m_c^5} \{ \langle \mathcal{O}_8^{\psi'}(^1S_0) \rangle + \frac{7}{m_c^2} \langle \mathcal{O}_8^{\psi'}(^3P_0) \rangle \} g(x_a, Q^2) g(x_b, Q^2) \right. \\ & \left. + \frac{\pi^3 \alpha_s^2}{54m_c^5} \langle \mathcal{O}_8^{\psi'}(^3S_1) \rangle \{ q(x_a, Q^2) \bar{q}(x_b, Q^2) + q \leftrightarrow \bar{q} \} \right], \quad (3) \end{aligned}$$

where x_a and x_b are the momentum fraction in a proton and given as

$$x_a = \frac{x_L + \sqrt{x_L^2 + 4\tau}}{2}, \quad x_b = \frac{-x_L + \sqrt{x_L^2 + 4\tau}}{2}, \quad x_L \equiv \frac{2p_L}{\sqrt{s}}, \quad \tau \equiv \frac{M^2}{s}, \quad (4)$$

with longitudinal momentum p_L of the produced particle. $\langle \mathcal{O}_8^{\psi'}(^1S_0) \rangle$, $\langle \mathcal{O}_8^{\psi'}(^3S_1) \rangle$ and $\langle \mathcal{O}_8^{\psi'}(^3P_0) \rangle$ are nonperturbative long–distance factors associated with the production of a $c\bar{c}$ pair in a color–octet 1S_0 , 3S_1 and 3P_0 states, respectively. From recent analysis for charmonium hadroproductions, the value of $\langle \mathcal{O}_8^{\psi'}(^3S_1) \rangle$ and of the combination are given as $\langle \mathcal{O}_8^{\psi'}(^3S_1) \rangle \sim$

$4.6 \times 10^{-3} [\text{GeV}^3]$, $\langle \mathcal{O}_8^{\psi'}(^1S_0) \rangle + \frac{7}{m_c^2} \langle \mathcal{O}_8^{\psi'}(^3P_0) \rangle \sim 5.2 \times 10^{-3} [\text{GeV}^3]$ [5], and also another combination $\frac{1}{3} \langle \mathcal{O}_8^{\psi'}(^1S_0) \rangle + \frac{1}{m_c^2} \langle \mathcal{O}_8^{\psi'}(^3P_0) \rangle \sim (5.9 \pm 1.9) \times 10^{-3} [\text{GeV}^3]$ [6]. Then we find the ratio as $\frac{\tilde{\Theta}}{\Theta} \equiv \frac{\langle \mathcal{O}_8^{\psi'}(^1S_0) \rangle - \frac{1}{m_c^2} \langle \mathcal{O}_8^{\psi'}(^3P_0) \rangle}{\langle \mathcal{O}_8^{\psi'}(^1S_0) \rangle + \frac{7}{m_c^2} \langle \mathcal{O}_8^{\psi'}(^3P_0) \rangle} \sim 0.69 \div 1.63$.

In order to study how the two-spin asymmetry $A_{LL}^{\psi'}$ is affected by the spin-dependent gluon distribution $\Delta g(x)$, we take a model of $\Delta g(x)$. So far, many people have suggested various kinds of $\Delta g(x)$ from the analysis of the data on $g_1(x, Q^2)$ [7, 8, 9]. The x -dependence of $x\Delta g(x, Q^2)$ at $Q^2 = 10\text{GeV}^2$ is shown in fig.1. We estimate the $A_{LL}^{\psi'}(pp)$ as a function of x_L for ψ' -productions at relevant RHIC energies, $\sqrt{s} = 50$ and 500GeV , by using these spin-dependent gluon distributions together with the spin-independent parton distribution of the Owens parametrization [10] for (a), the BBS parametrization [8] for (b), and the GRV92 LO parametrization [11] for (c), and taking Q^2 as $4m_c^2$ with $m_c = 1.5\text{GeV}$. Putting the ratio $\tilde{\Theta}/\Theta = 1$, we show the results of $A_{LL}^{\psi'}(pp)$ at $\sqrt{s} = 50$ and 500GeV in figs.2 and 3, respectively. For $\sqrt{s} = 50\text{GeV}$, we find that $A_{LL}^{\psi'}(pp)$ is very effective for examining not only the magnitude but also the behavior of the gluon polarization. But at $\sqrt{s} = 500\text{GeV}$ we need rather precise data on $A_{LL}^{\psi'}(pp)$ to test $\Delta g(x)$. There is a tendency that the smaller the $\sqrt{s}$ is, the larger the $A_{LL}^{\psi'}(pp)$ become. This is due to the fact that at larger $\sqrt{s}$, x_a and $x_b (= x_a - x_L)$ defined by eq.(4) take smaller value, and so $g(x_a)$ and $g(x_b)$ at $\sqrt{s} = 500\text{GeV}$ become large. Accordingly, $\Delta g(x_a)/g(x_a) \times \Delta g(x_b)/g(x_b)$ at $\sqrt{s} = 500\text{GeV}$ become smaller than that at $\sqrt{s} = 50\text{GeV}$.

In summary, observation of $A_{LL}^{\psi'}(pp)$ at small- p_T regions can largely contribute for the confirmation of the color-octet mechanism. At present, we do not know the exact value of $\tilde{\Theta}/\Theta$ which ranges in $0.69 \div 1.63$. It is interesting to fix this value from other experiments in order to make a precise prediction of $A_{LL}^{\psi'}(pp)$. Furthermore, the small- p_T ψ' -production allow us to give a rather clean test as a probe of the magnitude and x -dependence of the gluon polarization.

References

- [1] F. Abe et. al., CDF Collab., Phys. Rev. Lett. **69** (1992) 3704; Phys. Rev. Lett. **71** (1993) 2537.
- [2] E. Braaten and T. C. Yuan, Phys. Rev. Lett. **71** (1993) 1673; P. Cho and A. K. Leibovich, Phys. Rev. **D53** (1996) 150; W. -K. Tang and M. Vanttinen, Phys. Rev. **D53** (1996)

- 4851; S. Gupta and P. Mathews, preprint TIFR/TH/96-53, hep-ph/9609504 (1996).
- [3] M. Cacciari and M. Krämer, Phys. Rev. Lett. **76** (1996) 4128.
- [4] J. Ashman et al., EMC Collab., Phys. Lett. **B206** (1988) 364; Nucl. Phys. **B328** (1989) 1; B. Adeva et al., SMC Collab., Phys. Lett. **B302** (1993) 533; **B320** (1994) 400; P. L. Anthony et al., E142 Collab., Phys. Rev. Lett. **71** (1993) 959; D. Adeva et al., SMC Collab., Phys. Lett. **B329** (1994) 399; K. Abe et al., E143 Collab., Phys. Rev. Lett. **74** (1995) 346.
- [5] M. Beneke and I. Z. Rothstein, Phys. Rev. **D54** (1996) 2005.
- [6] M. Vanttinen, preprint CALT-68-2082, hep-ph/9610381 (1996).
- [7] T. Gehrmann and W. J. Stirling, Z. Phys. **C65** (1995) 461.
- [8] S. J. Brodsky, M. Burkardt and I. Schmidt, Nucl. Phys. **B441**, (1995) 197.
- [9] M. Glück, E. Reya and W. Vogelsang, Phys. Lett. **B359** (1995) 201.
- [10] J. F. Owens, Phys. Lett. **B266** (1991) 126.
- [11] M. Glück, E. Reya and W. Vogelsang, Z. Phys. **C53** (1992) 127.

Figure captions

Fig. 1: The x dependence of $x\Delta g(x, Q^2)$ at $Q^2 = 10\text{GeV}^2$ for various types of spin-dependent gluon distributions. The solid, dash-dotted, dotted, dashed and broken lines indicate the set A, B, C of ref.[7], ref.[8] and the ‘standard senrio’ of ref.[9], respectively.

Fig. 2: The two-spin asymmetry $A_{LL}^{\psi'}(pp)$ for $\tilde{\Theta}/\Theta = 1$ and $\sqrt{s} = 50\text{GeV}$, calculated with various types of $\Delta g(x)$, as a function of longitudinal momentum fraction x_L of ψ' . Various lines represent the same as in fig.1.

Fig. 3: The two-spin asymmetry $A_{LL}^{\psi'}(pp)$ for $\tilde{\Theta}/\Theta = 1$ and $\sqrt{s} = 500\text{GeV}$. Various lines represent the same as in fig.1.




Fig.1




Fig.2




Fig.3